



MEETING ABSTRACT

Open Access

Smoking alters the antigenicity and infectivity of *Porphyromonas gingivalis*

Iris Zeller¹, Justin A Hutcherson¹, Richard J Lamont^{1,2}, Donald R Demuth¹, Pinar Gumus³, Nejat Nizam³, Nurcan Buduneli³, David A Scott^{1*}

From 11th Annual Conference of the International Society for the Prevention of Tobacco Induced Diseases (ISPTID)

Athens, Greece. 9-11 December 2013

Background

Cigarette smokers are more susceptible to periodontal diseases and are more likely to be infected with *Porphyromonas gingivalis* than non-smokers. Furthermore, smoking is known to alter the expression of *P. gingivalis* surface components and to compromise IgG generation. The aim of this study was to evaluate if the overall IgG response to *P. gingivalis* is suppressed in smokers in vivo and if previously established in vitro tobacco-induced phenotypic *P. gingivalis* changes would be reflected in vivo.

Materials and methods

We examined the humoral response to several *P. gingivalis* strains as well as specific tobacco-regulated outer membrane proteins (FimA and RagB) by ELISA in biochemically-validated (salivary cotinine) smokers and non-smokers with chronic (CP, n = 13) or aggressive (AP, n = 20) periodontitis. We also monitored the local and systemic presence of *P. gingivalis* DNA by PCR.

Results

Smoking was associated with decreased total IgG responses against clinical (10512, 5607, and 10208C; all p < 0.05) but not laboratory (ATCC 33277, W83) *P. gingivalis* strains. Smoking did not influence IgG produced against specific cell surface proteins, although a non-significant pattern towards increased total FimA-specific IgG in CP subjects, but not AP subjects, was observed. Seropositive smokers were more likely to be

infected orally and systemically with *P. gingivalis* (p < 0.001), as determined by 16S RNA analysis.

Conclusions

Smoking alters the humoral response against *P. gingivalis* and may increase *P. gingivalis* infectivity, strengthening the evidence that mechanisms of periodontal disease progression in smokers may differ from non-smokers with the same disease classification.

Acknowledgements

This study was funded by NIDCR (R01DE019826 to DAS).

Authors' details

¹Oral Health and Systemic Disease, University of Louisville, Louisville, Kentucky, 40292, USA. ²School of Dentistry, University of Louisville, Louisville, Kentucky, 40292, USA. ³Department of Periodontology, School of Dentistry, Ege University, Izmir, 35040, Turkey.

Published: 6 June 2014

doi:10.1186/1617-9625-12-S1-A14

Cite this article as: Zeller et al.: Smoking alters the antigenicity and infectivity of *Porphyromonas gingivalis*. *Tobacco Induced Diseases* 2014 12(Suppl 1):A14.

* Correspondence: dascot07@louisville.edu

¹Oral Health and Systemic Disease, University of Louisville, Louisville, Kentucky, 40292, USA

Full list of author information is available at the end of the article